

考試科目	開課系級	考試日期	印製份數	答案紙	命題教師	備註
工程數學一	二 AB	1月19日		<input checked="" type="checkbox"/> 需 <input type="checkbox"/> 不需	陳桂鴻 呂學育	期末考

學生可帶 書本 計算機 其他_____ 皆不可

共 2 頁，第 1 頁

1. $A = \begin{bmatrix} 0 & 1 \\ -1 & 2 \end{bmatrix}$.

- (1) Find all eigenvalues and corresponding eigenvector. 3%
- (2) Find generalized eigenvector and obtain the transition matrix P of A . 5%
- (3) Find P^{-1} . 2%
- (4) Find the Jordan canonical form of A by using the similar transform ($P^{-1}AP$). 5%

2. $A = \begin{bmatrix} 1 & -2 \\ -2 & 1 \end{bmatrix}$

- (1) Find eigenvectors and write the transition matrix P of A . 3%
- (2) Find P^{-1} by using the orthogonal matrix property. 3%
- (3) Find the diagonal form of A by using the similar transform ($P^{-1}AP$). 3%
- (4) If $f(x) = x^{100}$, find the matrix $f(A)$ by using (a) the method of similar transform (matrix function), 9%
(b) Cayley-Hamilton theory. 9%
- (5) Find A^{-1} by using (a) adjoint method, 2% (b) Cayley-Hamilton theory. 6%

3. For the given linear system

$$-x_1 + 3x_2 = 0$$

$$x_1 - 2x_2 + x_3 = 1$$

$$x_2 + 2x_3 = 0$$

we can rewrite it as a matrix-vector equation $AX = B$

with the matrix $A = \begin{pmatrix} -1 & 3 & 0 \\ 1 & -2 & 1 \\ 0 & 1 & 2 \end{pmatrix}$, the vector $X = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$

- (1) write out the vector B (1%)
- (2) calculate $\det A$ (1%)
- (3) is the matrix A nonsingular? (1%)
- (4) write out the adjoint of the matrix A (2%)
- (5) find the inverse of the matrix A (2%)
- (6) solve the system to give the vector X (2%)
- (7) for the matrix A , what is the maximum number of independent column vectors? (1%)
- (8) what is the rank of the matrix A ? (2%)

4. For a given matrix $A = \begin{pmatrix} 0 & -1 & -1 \\ -1 & 0 & -1 \\ -1 & -1 & 0 \end{pmatrix}$

- (1) find the eigenvalues (hint: with 1 as an eigenvalue of multiplicity 2) (2%)
- (2) compute A^m ; $m = 10$ by using Cayley-Hamilton theorem (9%)
- (3) find a set of three mutually orthogonal eigenvectors (9%)
- (4) use these vectors obtained in (3) to construct an orthogonal matrix that diagonalizes the matrix A (3%)
- (5) compute A^m ; $m = 10$ by diagonalizing the matrix A (5%)

5. For a given conic section of the form $2xy = 1$

- (1) write the equation as the matrix product $X^T AX = 1$, with $X = \begin{pmatrix} x \\ y \end{pmatrix}$ (2%)
- (2) eliminate the xy -term by means of an orthogonal matrix and diagonalization (8%)