

海洋大學河海工程學系 2005 工程數學(四)期末考(Open Book)

1. Find the possible functions  $X_n(x)$  such that

$$\begin{aligned} X_n''(x) &= -\lambda X_n(x), \\ X_n'(0) &= X_n'(\pi) = 0 \end{aligned}$$

Also, find the eigenvalues  $\lambda_n$ . (20%)

2. A free-free string with a length  $\pi$

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial t^2}$$

$$u(x,0) = \cos(3x)$$

$$\dot{u}(x,0) = 0$$

$$u_x(0,t) = u_x(\pi,t) = 0$$

find  $u(\frac{\pi}{2}, \pi) = ?$

Using (1) Diamond rule (10%)

(2) Image method (10%)

(3) Series solution (10%)

3. Explain: (1) Laplace equation (5%)

(2) Wave equation (5%)

(3) Heat equation (5%)

(4) Characteristic line (5%)

(5) D'Alembert solution (5%)

4. Solve the PDE:  $u_{tt} = \begin{cases} 4u_{xx}, & x < 0, t > 0 \\ 1u_{xx}, & x > 0, t > 0 \end{cases}$

$$\text{I.C.: } u(x,0) = \dot{u}(x,0) = 0$$

At the interface, we apply the force

$$u_x(0^+, t) - u_x(0^-, t) = a \sin \omega t. \quad (20\%)$$

Hint: Using diamond rule




5. Solve  $u_{tt} = u_{xx}$

$$u(x,0) = 0$$

$$\dot{u}(x,0) = \frac{1}{a} [H(x+a) - H(x-a)]$$

(1) Solve  $u(x,t)$  for  $a=1$ . (10%)

(2) Solve  $u(x,t)$  for  $a \rightarrow 0$ . (10%)



$$H(x) = \begin{cases} 1, & x > 0 \\ 0, & x < 0 \end{cases}$$