

Dear Author,

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For fax submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/ corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

#### **Please note**

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL: [http://dx.doi.org/\[DOI\]](http://dx.doi.org/[DOI]).

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information go to: <http://www.springerlink.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us if you would like to have these documents returned.

# Metadata of the article that will be visualized in OnlineFirst

---

ArticleTitle	New series expansions for fundamental solutions of linear elastostatics in 2D	
--------------	---	--

---

Article Sub-Title		
-------------------	--	--

---

Article CopyRight	Springer-Verlag (This will be the copyright line in the final PDF)	
-------------------	---	--

---

Journal Name	Computing	
--------------	-----------	--

---

Corresponding Author	Family Name	<b>Lee</b>
	Particle	
	Given Name	<b>Ming-Gong</b>
	Suffix	
	Division	Department of Applied Statistics
	Organization	Chung Hua University
	Address	Hsin-Chu, Taiwan
	Email	mglee@chu.edu.tw

---

Author	Family Name	<b>Li</b>
	Particle	
	Given Name	<b>Zi-Cai</b>
	Suffix	
	Division	Department of Applied Mathematics
	Organization	National Sun Yat-sen University
	Address	Kaohsiung, 80424, Taiwan
	Division	Department of Computer Science and Engineering
	Organization	National Sun Yat-sen University
	Address	Kaohsiung, 80424, Taiwan
	Division	Department of Applied Mathematics
	Organization	Chung Hua University
	Address	Hsin-Chu, Taiwan
	Email	zcli@math.nsysu.edu.tw

---

Author	Family Name	<b>Chen</b>
	Particle	
	Given Name	<b>Jeng-Tzong</b>
	Suffix	
	Division	Department of Harbor and River Engineering
	Organization	National Taiwan Ocean University
	Address	Keelung, 20224, Taiwan
	Email	jtchen@mail.ntou.edu.tw

---

Schedule	Received	02 July 2010
	Revised	
	Accepted	28 October 2010

---

Abstract	Series expansions of fundamental solutions are essential to algorithms and analysis of the null field method (NFM) and to analysis of the method of fundamental solutions (MFS). For linear elastostatics, new Fourier series expansions of FS are derived, directly from integration. The new expansions of the FS are simpler than	
----------	--	--

those in Chen et al. [J Mech 26(3):113–121, 2010], thus facile to application in NFM and MFS. The new series expansions of FS in this paper are important to both theory and computation of linear elastostatics.

---

Keywords (separated by '-') Elastostatics - Fundamental solutions - Expansions of fundamental solutions - Method of fundamental solutions - Null field method

---

Footnote Information

---

Journal: 607  
Article: 137



## Author Query Form

**Please ensure you fill out your response to the queries raised below and return this form along with your corrections**

Dear Author

During the process of typesetting your article, the following queries have arisen. Please check your typeset proof carefully against the queries listed below and mark the necessary changes either directly on the proof/online grid or in the ‘Author’s response’ area provided below

Query	Details required	Author’s response
1.	Please check and confirm the author names and initials are correct. Also, kindly confirm the details in the metadata are correct.	
2.	Please check the “)” has been deleted in Equation 2.15	
3.	Please check the breaking of the equation 5.17.	
4.	Please provide MSC codes. For more details, if required, kindly visit <a href="http://www.ams.org/msc/">http://www.ams.org/msc/</a> .	

# New series expansions for fundamental solutions of linear elastostatics in 2D

Zi-Cai Li · Ming-Gong Lee · Jeng-Tzong Chen

Received: 2 July 2010 / Accepted: 28 October 2010  
© Springer-Verlag 2010

**Abstract** Series expansions of fundamental solutions are essential to algorithms and analysis of the null field method (NFM) and to analysis of the method of fundamental solutions (MFS). For linear elastostatics, new Fourier series expansions of FS are derived, directly from integration. The new expansions of the FS are simpler than those in Chen et al. [J Mech 26(3):113–121, 2010], thus facile to application in NFM and MFS. The new series expansions of FS in this paper are important to both theory and computation of linear elastostatics.

**Keywords** Elastostatics · Fundamental solutions · Expansions of fundamental solutions · Method of fundamental solutions · Null field method

---

Z.-C. Li  
Department of Applied Mathematics, National Sun Yat-sen University,  
Kaohsiung 80424, Taiwan  
e-mail: zcli@math.nsysu.edu.tw

Z.-C. Li  
Department of Computer Science and Engineering, National Sun Yat-sen University,  
Kaohsiung 80424, Taiwan

Z.-C. Li  
Department of Applied Mathematics, Chung Hua University, Hsin-Chu, Taiwan

M.-G. Lee (✉)  
Department of Applied Statistics, Chung Hua University, Hsin-Chu, Taiwan  
e-mail: mglee@chu.edu.tw

J.-T. Chen  
Department of Harbor and River Engineering, National Taiwan Ocean University,  
Keelung 20224, Taiwan  
e-mail: jtchen@mail.ntou.edu.tw

## 1 Introduction

Series expansions of fundamental solutions are essential to error and stability analysis of the method of fundamental solutions, and to algorithms and analysis of the null field method (NFM). For linear elastostatics, new Fourier series expansions of FS are derived, directly from integration. The new expansions of the FS are simpler than those in Chen et al. [5], thus facile to analysis of MFS [12–14] and to the NFM [16–18]. For elasticity problems, when the fundamental solutions (FS) or particular solutions (PS) satisfying PDE are chosen, the Trefftz method (TM) [19] leads to the method of fundamental solutions (MFS) and the method of particular method (MPS), respectively. The MFS and MPS can be applied to arbitrary domains. For circular domain with multiple holes, the null field method (NFM) is proposed by Chen with his research groups. In NFM, the fundamental solutions (FS) with the source nodes outside of the solution domain are used in the Green formulas. The Fourier expansions of the known and the unknown boundary conditions on the circular boundaries are chosen, so that the explicit algebraic equations are easily obtained by means of orthogonality of Fourier functions. Recently, the explicit linear algebraic equations of NFM are provided in [16–18], which are easy for real application. The NFM has been applied to elliptic and eigenvalue problems in circular domains with multiple holes, reported in many papers; here we cite Chen's work for Laplace's equation in [2, 4], and for elastostatics problems in [3, 5]. Extensions to elliptic boundaries can be found in Crouch and Mogilevskaya [8], and Chen et al. [6, 7].

For the NFM of elastostatics problems, the expansions of the fundamental solutions (FS) are a must. Although the series expansions of FS for plane elastostatics and basic methods are first given in Chen et al. [5]. In this paper, new and simpler series expansions of the FS are derived directly via integration. On the other hand, for error analysis of the MFS, the expansions of FS are essential. Since the error bounds of harmonic polynomials as PS in TM have been established in [19], the errors between the FS and harmonic polynomials can be found, based on the series expansions of FS, see Li [12] for Laplace's equation. The stability analysis of MFS in [15] also needs the expansions of FS. By following the arguments in [12, 15], we may also carry out the analysis for the MFS of linear elastostatics (see Li [14]), once the series expansions of FS are provided. Hence, the new series expansions of FS in this paper are *important to both theory and computation of linear elastostatics* [10, 11, 19–22].

This paper is organized as follows. In Sect. 2, basic mathematical description for linear elastostatics problems in 2D is provided, and their fundamental solutions (FS) are introduced. In Sect. 3, preliminary expansion formulas are given, and in Sects. 4 and 5 main integration formula are derived. In Sect. 6, new expansions of the FS of linear elastostatics in 2D are provided.

## 2 Linear elastostatics problems in 2D

### 2.1 Basic theory

Consider the linear elastostatics problem in 2D. Denote the displacement vector,

$$\vec{w} = \mathbf{w} = \{w_1(\mathbf{x}), w_2(\mathbf{x})\}^T = \{u(x, y), v(x, y)\}^T, \quad (2.1)$$

52 where  $\vec{x} = \mathbf{x} = (x_1, x_2) = (x, y)$ . When there exists no body force  $\vec{f} \equiv 0$ , we obtain  
 53 the homogeneous equation, called the Cauchy–Navier equation of linear elastostatics  
 54 for isotropic body:

$$\mu \Delta \vec{w} + (\lambda + \mu) \nabla(\nabla \cdot \vec{w}) = 0 \quad \text{in } S, \tag{2.2}$$

56 where  $\lambda$  and  $\mu$  are the Lamé constants. Eq. (2.2) can be written as

$$\Delta \vec{w} + \frac{1}{1 - 2\nu} \nabla(\nabla \cdot \vec{w}) = 0 \quad \text{in } S, \tag{2.3}$$

58 where the Poisson ratio

$$\nu = \frac{\lambda}{2(\lambda + \mu)}, \quad 0 < \nu < \frac{1}{2}. \tag{2.4}$$

60 We cite a theorem from Chen and Zhou [1].

61 **Theorem 2.1** *The general solutions of the linear elastostatic equations (2.2) in 3D*  
 62 *and 2D are given by*

$$\vec{w}(\vec{x}) = \vec{h}(\vec{x}) - \kappa \nabla[\vec{x} \cdot \vec{h}(\vec{x}) + q(\vec{x})], \tag{2.5}$$

64 where  $\vec{h}(\vec{x})$  are the harmonic vector and  $q(\vec{x})$  is a harmonic function, and the constant

$$\kappa = \frac{1}{4(1 - \nu)}. \tag{2.6}$$

## 66 2.2 Fundamental solutions

67 The principal fundamental solutions of linear Elastostatics in 2D are given in [1] as

$$E_2(\mathbf{x}, \xi) = \frac{\lambda + 3\mu}{4\pi\mu(\lambda + 2\mu)} \left\{ -\ln r_{\mathbf{x}\xi} I_2 + \frac{\lambda + \mu}{\lambda + 3\mu} \frac{1}{r_{\mathbf{x}\xi}^2} [(\mathbf{x} - \xi)(\mathbf{x} - \xi)^T] \right\}, \tag{2.7}$$

69 where  $r_{\mathbf{x}\xi} = |\mathbf{x} - \xi|$  and  $I_2$  is the identity matrix. The FS in (2.7) satisfy

$$\mu \Delta E_2(\mathbf{x}, \xi) + (\lambda + \mu) \nabla_{\mathbf{x}}[\nabla_{\mathbf{x}} \cdot E_2(\mathbf{x}, \xi)] = -\delta(\mathbf{x} - \xi) I_2. \tag{2.8}$$

71 Choose the source nodes  $Q(\xi_i, \eta_i)$  to be uniformly located on a larger circle of the  
 72 2D domain  $S$ , and denote the collocation points  $P(x, y)$ , where

$$\begin{aligned} x &= \rho \cos \theta, & y &= \rho \sin \theta, \\ \xi_i &= R \cos \phi_i, & \eta_i &= R \sin \phi_i, \quad i = 1, 2, \dots, N, \end{aligned} \tag{2.9}$$

75 where  $R > \max_S \rho$ ,  $\rho = \sqrt{x^2 + y^2}$ ,  $R = \sqrt{\xi_i^2 + \eta_i^2}$  and  $\phi_i = \frac{i2\pi}{N}$ . Then

$$76 \quad r_i = r_{x\xi_i} = \sqrt{R^2 + \rho^2 - 2R\rho \cos(\theta - \phi_i)}. \quad (2.10)$$

77 Denote  $\vec{d}_i = (a_i, b_i)$  with the constants  $a_i$  and  $b_i$ , and  $\vec{u}_i = (u_i, v_i)^T$ . We have

$$78 \quad \vec{u}_i = E_2(\mathbf{x}, \xi_i)\vec{d}_i, \quad (2.11)$$

79 where

$$80 \quad u_i = a_i \left( -A \ln r_i + B \frac{(x - \xi_i)^2}{r_i^2} \right) + b_i B \frac{(x - \xi_i)(y - \eta_i)}{r_i^2}, \quad (2.12)$$

$$81 \quad v_i = a_i B \frac{(x - \xi_i)(y - \eta_i)}{r_i^2} + b_i \left( -A \ln r_i + B \frac{(y - \eta_i)^2}{r_i^2} \right), \quad (2.13)$$

82 and the constants

$$83 \quad A = \frac{\lambda + 3\mu}{4\pi\mu(\lambda + 2\mu)}, \quad B = \frac{\lambda + \mu}{4\pi\mu(\lambda + 2\mu)}. \quad (2.14)$$

84 From Theorem 2.1, by adding the simple fundamental solutions  $\nabla(\ln r) = \left\{ \frac{x}{r^2}, \frac{y}{r^2} \right\}^T$ ,  
85 we obtain the general linear combination of fundamental solutions,

$$86 \quad u_N = u_N(x, y) = \sum_{i=1}^N \left\{ a_i \left( -\ln r_i + D \frac{(x - \xi_i)^2}{r_i^2} \right) + b_i D \frac{(x - \xi_i)(y - \eta_i)}{r_i^2} \right. \\ 87 \quad \left. + c_i \frac{(x - \xi_i)}{r_i^2} \right\}, \quad (2.15)$$

$$88 \quad v_N = v_N(x, y) = \sum_{i=1}^N \left\{ a_i D \frac{(x - \xi_i)(y - \eta_i)}{r_i^2} + b_i \left( -\ln r_i + D \frac{(y - \eta_i)^2}{r_i^2} \right) \right. \\ 89 \quad \left. + c_i \frac{(y - \eta_i)}{r_i^2} \right\}, \quad (2.16)$$

90 where  $a_i$ ,  $b_i$  and  $c_i$  are the coefficients, the constant

$$91 \quad D = \frac{B}{A} = \frac{\lambda + \mu}{\lambda + 3\mu} = \frac{1}{3 - 4\nu} = \frac{\kappa}{1 - \kappa}, \quad (2.17)$$

92 and  $\kappa$  is given in (2.6). By using the FS in (2.15) and (2.16), and by employing the  
93 collocation Trefftz techniques in [19], the method of fundamental solutions (MFS)



94 are established. For Laplace’s equation, the error analysis of MFS in [14] is made,  
 95 based on the expansions of FS (see Li [12]), and the stability analysis of MFS in [15]  
 96 also needs the expansions of FS. Evidently, the expansions of FS are a basic tool in  
 97 analysis of the MFS.

98 The principal FS in (2.7) is expressed in the matrix form

$$\begin{aligned}
 & \begin{pmatrix} -\ln r(1-\kappa) + \kappa \frac{x^2}{r^2} & \kappa \frac{xy}{r^2} \\ \kappa \frac{xy}{r^2} & -\ln r(1-\kappa) + \kappa \frac{y^2}{r^2} \end{pmatrix} \\
 & = (1-\kappa) \begin{pmatrix} -\ln r + D \frac{x^2}{r^2} & D \frac{xy}{r^2} \\ D \frac{xy}{r^2} & -\ln r + D \frac{y^2}{r^2} \end{pmatrix}, \tag{2.18}
 \end{aligned}$$

101 where  $D$  is given in (2.17). We cite the Betti–Somigliana formula with the principal  
 102 FS [1].

103 **Theorem 2.2** *There exist the Betti–Somigliana formulas for linear elastostatics:*

$$\bar{w}(\mathbf{x}) = \int_{\partial\Omega} \{E_2(\mathbf{x}, \xi) \cdot \bar{\tau}(\bar{w}) - \bar{\tau}(E_2(\mathbf{x}, \xi)) \cdot \bar{w}\} d\sigma_\xi, \quad \mathbf{x} \in \Omega, \tag{2.19}$$

$$\frac{1}{2} \bar{w}(\mathbf{x}) = \int_{\partial\Omega} \{E_2(\mathbf{x}, \xi) \cdot \bar{\tau}(\bar{w}) - \bar{\tau}(E_2(\mathbf{x}, \xi)) \cdot \bar{w}\} d\sigma_\xi, \quad \mathbf{x} \in \partial\Omega, \tag{2.20}$$

$$0 = \int_{\partial\Omega} \{E_2(\mathbf{x}, \xi) \cdot \bar{\tau}(\bar{w}) - \bar{\tau}(E_2(\mathbf{x}, \xi)) \cdot \bar{w}\} d\sigma_\xi, \quad \mathbf{x} \in \bar{\Omega}^c, \tag{2.21}$$

107 where  $\Omega$  is an open area, and  $\bar{\Omega}^c$  is the complementary region of the close area  $\bar{\Omega}$   
 108 including the exterior and the interior boundaries  $\partial S_R$  and  $\partial S_{R_1}$ .

109 The boundary elements methods originated from (2.20) and (2.19), but the NFM is  
 110 developed from (2.21) and (2.19), based on the expansions of FS (see [5]).

### 111 3 Preliminary formulas

112 Denote two nodes by  $P(x, y)$  and  $Q(\xi, \eta)$  with  $x = \rho \cos \theta, y = \rho \sin \theta, \xi =$   
 113  $R \cos \phi,$  and  $\eta = R \sin \phi.$  Then  $\rho = \sqrt{x^2 + y^2}, R = \sqrt{\xi^2 + \eta^2}$  and  $r = |\overline{PQ}| =$   
 114  $\sqrt{\rho^2 - 2\rho R \cos(\theta - \phi) + R^2}.$  In this section, we give the basic integral expansions  
 115 of  $\ln r$  and  $\frac{1}{r^2}.$  First we obtain a lemma from [9, 12].

116 **Lemma 3.1** For  $R > \rho$ , there exist the equalities

117 
$$\frac{1}{2\pi} \int_0^{2\pi} (\ln \sqrt{\rho^2 - 2\rho R \cos(\theta - \phi) + R^2}) d\phi = \ln R,$$

118 
$$\frac{1}{\pi} \int_0^{2\pi} (\ln \sqrt{\rho^2 - 2\rho R \cos(\theta - \phi) + R^2}) \cos m\phi d\phi = -\frac{\rho^m}{mR^m} \cos m\theta, \quad m = 1, 2, \dots,$$

119 
$$\frac{1}{\pi} \int_0^{2\pi} (\ln \sqrt{\rho^2 - 2\rho R \cos(\theta - \phi) + R^2}) \sin m\phi d\phi = -\frac{\rho^m}{mR^m} \sin m\theta, \quad m = 1, 2, \dots$$

120 Based on the Fourier coefficients of  $\ln \sqrt{\rho^2 - 2\rho R \cos(\theta - \phi) + R^2}$  from  
121 Lemma 3.1, we have the following lemma immediately.

122 **Lemma 3.2** There exist the expansions

123 
$$\ln \sqrt{\rho^2 - 2\rho R \cos(\theta - \phi) + R^2} = \ln R - \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{\rho}{R}\right)^n \cos n(\theta - \phi), \quad \rho < R, \quad (3.1)$$

124 
$$\ln \sqrt{\rho^2 - 2\rho R \cos(\theta - \phi) + R^2} = \ln \rho - \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{R}{\rho}\right)^n \cos n(\theta - \phi), \quad \rho > R. \quad (3.2)$$

125 Equations (3.1) and (3.2) are the expansions of the FS for Laplace's equation in  
126 2D used in [12, 15]. The rest of this paper is devoted to derive new expansions for  
127  $\kappa \nabla(\ln r) = \kappa \left\{ \frac{x}{r^2}, \frac{y}{r^2} \right\}^T$  and (2.18) of linear Elastostatics in 2D. We give a new lemma.

128 **Lemma 3.3** There exist the integrals,

129 
$$\int_0^{2\pi} \frac{\cos nx}{1 + a^2 - 2a \cos x} dx = \frac{2\pi a^n}{1 - a^2} (a^2 < 1) = \frac{2\pi a^{-n}}{a^2 - 1} (a^2 > 1), \quad (3.3)$$

130 
$$\int_0^{2\pi} \frac{\sin nx}{1 + a^2 - 2a \cos x} dx = 0. \quad (3.4)$$

131 *Proof* From Gradsheyan and Ryzhik, [9], p. 366

132 
$$\int_0^{\pi} \frac{\cos nx}{1 + a^2 - 2a \cos x} dx = \frac{\pi a^n}{1 - a^2} (a^2 < 1) = \frac{\pi a^{-n}}{a^2 - 1} (a^2 > 1), \quad (3.5)$$

133 we have

$$\begin{aligned}
 134 \quad \int_0^{2\pi} \frac{\cos nx}{1+a^2-2a\cos x} dx &= \int_{-\pi}^{\pi} \frac{\cos nx}{1+a^2-2a\cos x} dx \\
 135 \quad &= \left( \int_{-\pi}^0 + \int_0^{\pi} \right) \frac{\cos nx}{1+a^2-2a\cos x} dx. \quad (3.6)
 \end{aligned}$$

136 Let  $x = -t$  and  $dx = -dt$ . We have

$$\begin{aligned}
 137 \quad \int_{-\pi}^0 \frac{\cos nx}{1+a^2-2a\cos x} dx &= - \int_{\pi}^0 \frac{\cos nx}{1+a^2-2a\cos x} dx \\
 138 \quad &= \int_0^{\pi} \frac{\cos nx}{1+a^2-2a\cos x} dx. \quad (3.7)
 \end{aligned}$$

139 Combining (3.5)–(3.7) gives

$$\begin{aligned}
 140 \quad \int_0^{2\pi} \frac{\cos nx}{1+a^2-2a\cos x} dx &= 2 \int_0^{\pi} \frac{\cos nx}{1+a^2-2a\cos x} dx \\
 141 \quad &= \frac{2\pi a^n}{1-a^2} (a^2 < 1) = \frac{2\pi a^n}{a^2-1} (a^2 > 1). \quad (3.8)
 \end{aligned}$$

142 This is the first desired result (3.3) with  $a^2 < 1$ . The proof for (3.3) with  $a^2 > 1$  is  
 143 similar.

144 Next we have

$$145 \quad \int_0^{2\pi} \frac{\sin nx}{1+a^2-2a\cos x} dx = \left( \int_{-\pi}^0 + \int_0^{\pi} \right) \frac{\sin nx}{1+a^2-2a\cos x} dx = 0, \quad (3.9)$$

146 since the following equality holds,

$$147 \quad \int_{-\pi}^0 \frac{\sin nx}{1+a^2-2a\cos x} dx = - \int_0^{\pi} \frac{\sin nx}{1+a^2-2a\cos x} dx = 0. \quad (3.10)$$

148 This is the second desired result (3.4), and completes the proof of Lemma 3.3.

149 **Lemma 3.4** For  $a^2 < 1$ , there exist the Fourier expansions,

$$150 \quad \frac{1}{1+a^2-2a\cos x} = \frac{2}{1-a^2} \left( \frac{1}{2} + \sum_{n=1}^{\infty} a^n \cos nx \right). \quad (3.11)$$

151 For  $a^2 > 1$ ,

$$152 \quad \frac{1}{1+a^2-2a\cos x} = \frac{2}{a^2-1} \left( \frac{1}{2} + \sum_{n=1}^{\infty} a^{-n} \cos nx \right). \quad (3.12)$$

153 *Proof* Denote the Fourier series

$$154 \quad \frac{1}{1+a^2-2a\cos x} = \frac{a_0}{2} + \sum_{k=1}^{\infty} (a_k \cos kx + b_k \sin kx), \quad (3.13)$$

155 where the Fourier coefficients  $a_k$  and  $b_k$  are obtained:

$$156 \quad a_k = \frac{1}{\pi} \int_0^{2\pi} \frac{\cos kx}{1+a^2-2a\cos x} dx, \quad k = 0, 1, 2, \dots, \quad (3.14)$$

$$157 \quad b_k = \frac{1}{\pi} \int_0^{2\pi} \frac{\sin kx}{1+a^2-2a\cos x} dx, \quad k = 1, 2, \dots \quad (3.15)$$

158 Then from Lemma 3.3 we obtain for  $a^2 < 1$

$$159 \quad a_k = \frac{2}{\pi} \frac{\pi a^k}{1-a^2} = \frac{2a^k}{1-a^2}, \quad k = 0, 1, \dots, \quad (3.16)$$

$$160 \quad b_k = 0, \quad k = 1, 2, \dots \quad (3.17)$$

161 Substituting the coefficients  $a_k$  and  $b_k$  into (3.13) gives the desired result (3.11) with  
 162  $a^2 < 1$ . Similarly, Eq. (3.12) with  $a^2 > 1$  also holds. This completes the proof of  
 163 Lemma 3.4.

164 **Theorem 3.1** There exist the expansions

$$165 \quad \frac{1}{r^2} = \frac{2}{R^2-\rho^2} \left( \frac{1}{2} + \sum_{n=1}^{\infty} \left( \frac{\rho}{R} \right)^n \cos n(\phi-\theta) \right), \quad \rho < R, \quad (3.18)$$

$$166 \quad \frac{1}{r^2} = \frac{2}{\rho^2-R^2} \left( \frac{1}{2} + \sum_{n=1}^{\infty} \left( \frac{R}{\rho} \right)^n \cos n(\phi-\theta) \right), \quad \rho > R, \quad (3.19)$$

167 where  $r^2 = R^2 + \rho^2 - 2\rho R \cos(\phi - \theta)$ ,  $\mathbf{x} = (x, y) = (\rho \cos \theta, \rho \sin \theta)$  and  $\xi =$   
 168  $(\xi, \eta) = (R \cos \phi, R \sin \phi)$ .

169 *Proof* Since  $r^2 = R^2(1 + a^2 - 2a \cos(\phi - \theta))$ , we obtain (3.18) from Lemma 3.4  
 170 by noting  $x = \phi - \theta$ . Eq. (3.19) follows from (3.18) by the symmetry between  $\rho$  and  
 171  $R$ . This completes the proof of Theorem 3.1.

172 **4 Expansions for  $\frac{x-\xi}{r^2}$  and  $\frac{y-\eta}{r^2}$**

173 The simple FS for  $q = \ln r$  is given from (2.5),

174 
$$\nabla q = \left( \frac{x - \xi}{r^2}, \frac{y - \eta}{r^2} \right)^T = (T_1, T_2)^T. \tag{4.1}$$

175 To obtain the Fourier expansions of (4.1), we need the following lemma.

176 **Lemma 4.1** *Let  $\rho < R$  and  $r^2 = R^2 + \rho^2 - 2\rho R \cos(\phi - \theta)$ . There exist the integral*  
 177 *formulas,*

178 
$$\int_0^{2\pi} \frac{x - \xi}{r^2} d\phi = \int_0^{2\pi} \frac{y - \eta}{r^2} d\phi = 0, \tag{4.2}$$

179 
$$\int_0^{2\pi} \frac{x - \xi}{r^2} \cos n\phi d\phi = -\frac{\pi}{R} \left(\frac{\rho}{R}\right)^{n-1} \cos(n - 1)\theta, \tag{4.3}$$

180 
$$\int_0^{2\pi} \frac{x - \xi}{r^2} \sin n\phi d\phi = -\frac{\pi}{R} \left(\frac{\rho}{R}\right)^{n-1} \sin(n - 1)\theta, \tag{4.4}$$

181 
$$\int_0^{2\pi} \frac{y - \eta}{r^2} \cos n\phi d\phi = \frac{\pi}{R} \left(\frac{\rho}{R}\right)^{n-1} \sin(n - 1)\theta, \tag{4.5}$$

182 
$$\int_0^{2\pi} \frac{y - \eta}{r^2} \sin n\phi d\phi = -\frac{\pi}{R} \left(\frac{\rho}{R}\right)^{n-1} \cos(n - 1)\theta. \tag{4.6}$$

183 *Proof* Denote  $a = \frac{\rho}{R}$ . We have from Theorem 3.1 and the orthogonality of trigono-  
 184 metric functions,

185 
$$\int_0^{2\pi} \frac{\rho \cos \theta - R \cos \phi}{r^2} d\phi = \rho \cos \theta \int_0^{2\pi} \frac{1}{\rho^2 + R^2 - 2\rho R \cos(\phi - \theta)} d\phi$$
  
 186 
$$- R \int_0^{2\pi} \frac{\cos \phi}{\rho^2 + R^2 - 2\rho R \cos(\phi - \theta)} d\phi$$
  
 187 
$$= \rho \cos \theta \frac{1}{R^2} \frac{2\pi}{1 - a^2} - \frac{1}{R^2} \frac{2\pi}{1 - a^2} R a \cos \theta = 0. \tag{4.7}$$

This is the left hand side of (4.2), and the proof for the right side of (4.2) is similar.

Next for (4.3), the integrand is given by

$$\begin{aligned}(x - \xi) \cos n\phi &= (\rho \cos \theta - R \cos \phi) \cos n\phi \\ &= \rho \cos \theta \cos n\phi - \frac{R}{2}(\cos(n-1)\phi + \cos(n+1)\phi).\end{aligned}\quad (4.8)$$

From Theorem 3.1 and the orthogonality of trigonometric functions,

$$\begin{aligned}\int_0^{2\pi} \frac{x - \xi}{r^2} \cos n\phi d\phi &= \frac{2\pi}{R^2(1-a^2)} \left\{ \rho \cos \theta a^n \cos n\theta \right. \\ &\quad \left. - \frac{R}{2}(a^{n-1} \cos(n-1)\theta + a^{n+1} \cos(n+1)\theta) \right\} \\ &= \frac{2\pi}{R^2(1-a^2)} \left\{ \frac{\rho a^n}{2}(\cos(n-1)\theta + \cos(n+1)\theta) \right. \\ &\quad \left. - \frac{R}{2}(a^{n-1} \cos(n-1)\theta + a^{n+1} \cos(n+1)\theta) \right\} \\ &= \frac{2\pi}{R^2(1-a^2)} a^{n-1} \left( \frac{\rho a}{2} - \frac{R}{2} \right) \cos(n-1)\theta.\end{aligned}\quad (4.9)$$

Since

$$\frac{\rho a}{2} - \frac{R}{2} = \frac{\rho^2}{2R} - \frac{R}{2} = \frac{1}{2R}(\rho^2 - R^2) = \frac{R}{2}(a^2 - 1),\quad (4.10)$$

the desired result (4.3) follows from (4.9).

Third for (4.6), the integrand is

$$\begin{aligned}(y - \eta) \sin n\phi &= (\rho \sin \theta - R \sin \phi) \sin n\phi \\ &= \rho \sin \theta \sin n\phi - \frac{R}{2}(\cos(n-1)\phi - \cos(n+1)\phi).\end{aligned}\quad (4.11)$$

Similarly, we have

$$\begin{aligned}\int_0^{2\pi} \frac{y - \eta}{r^2} \sin n\phi d\phi &= \frac{2\pi}{R^2(1-a^2)} \left\{ \rho \sin \theta a^n \sin n\theta \right. \\ &\quad \left. - \frac{R}{2}(a^{n-1} \cos(n-1)\theta - a^{n+1} \cos(n+1)\theta) \right\} \\ &= \frac{2\pi}{R^2(1-a^2)} \left\{ \frac{\rho a^n}{2}(\cos(n-1)\theta - \cos(n+1)\theta) \right. \\ &\quad \left. - \frac{R}{2}(a^{n-1} \cos(n-1)\theta - a^{n+1} \cos(n+1)\theta) \right\}\end{aligned}$$

$$\begin{aligned}
 &= \frac{2\pi}{R^2(1-a^2)} a^{n-1} \left( \frac{\rho a}{2} - \frac{R}{2} \right) \cos(n-1)\theta \\
 &= -\frac{\pi}{R} a^{n-1} \cos(n-1)\theta.
 \end{aligned} \tag{4.12}$$

This is the last result (4.6), and the proof for (4.4) and (4.5) is similar. This completes the proof of Lemma 4.1.

**Theorem 4.1** Let  $\rho < R$  and  $r^2 = R^2 + \rho^2 - 2\rho R \cos(\phi - \theta)$ . There exist the expansions<sup>1</sup>,

$$\begin{aligned}
 T_1^i &= \frac{x - \xi}{r^2} = -\frac{1}{R} \sum_{n=1}^{\infty} \left( \frac{\rho}{R} \right)^{n-1} (\cos(n-1)\theta \cos n\phi + \sin(n-1)\theta \sin n\phi) \\
 &= -\frac{1}{R} \sum_{n=0}^{\infty} \left( \frac{\rho}{R} \right)^n \cos(n(\theta - \phi) - \phi),
 \end{aligned} \tag{4.13}$$

$$\begin{aligned}
 T_2^i &= \frac{y - \eta}{r^2} = \frac{1}{R} \sum_{n=1}^{\infty} \left( \frac{\rho}{R} \right)^{n-1} (\sin(n-1)\theta \cos n\phi - \cos(n-1)\theta \sin n\phi) \\
 &= \frac{1}{R} \sum_{n=0}^{\infty} \left( \frac{\rho}{R} \right)^n \sin(n(\theta - \phi) - \phi).
 \end{aligned} \tag{4.14}$$

*Proof* We have the Fourier expansion,

$$\frac{x - \xi}{r^2} = \frac{\alpha_0}{2} + \sum_{k=1}^{\infty} (\alpha_k \cos k\phi + \beta_k \sin k\phi), \tag{4.15}$$

where

$$\alpha_k = \frac{1}{\pi} \int_0^{2\pi} \frac{x - \xi}{r^2} \cos k\phi, \quad \beta_k = \frac{1}{\pi} \int_0^{2\pi} \frac{x - \xi}{r^2} \sin k\phi. \tag{4.16}$$

From Lemma 4.1, we obtain the first equality of (4.13). Let  $n - 1 = k$  we have from (4.13)

$$\begin{aligned}
 \frac{x - \xi}{r^2} &= -\frac{1}{R} \sum_{k=1}^{\infty} \left( \frac{\rho}{R} \right)^k (\cos k\theta \cos(k+1)\phi + \sin k\theta \sin(k+1)\phi) \\
 &= -\frac{1}{R} \sum_{k=0}^{\infty} \left( \frac{\rho}{R} \right)^k \cos(k(\theta - \phi) - \phi).
 \end{aligned} \tag{4.17}$$

The proof for (4.14) is similar, and completes the proof of Theorem 4.1.

<sup>1</sup> The superscript “i” of  $T_1^i$  denotes the case of  $\rho < R$ , and the superscript “e” in Theorem 6.3 denotes the case of  $\rho > R$ .

228 **5 Expansions for  $\frac{(x-\xi)^2}{r^2}$ ,  $\frac{(y-\eta)^2}{r^2}$  and  $\frac{(x-\xi)(y-\eta)}{r^2}$**

229 For expansions of the principal FS, we need the following lemma.

230 **Lemma 5.1** *Let  $\rho < R$ ,  $a = \frac{\rho}{R}$  and  $r^2 = R^2 + \rho^2 - 2\rho R \cos(\phi - \theta)$ . There exist the*  
 231 *integral formulas*

232 
$$\int_0^{2\pi} \frac{(x-\xi)^2}{r^2} d\phi = \int_0^{2\pi} \frac{(y-\eta)^2}{r^2} d\phi = \pi, \quad (5.1)$$

233 
$$\int_0^{2\pi} \frac{(x-\xi)(y-\eta)}{r^2} d\phi = 0, \quad (5.2)$$

234 
$$\int_0^{2\pi} \frac{(x-\xi)^2}{r^2} \cos \phi d\phi = -\frac{\pi}{2} a \cos \theta, \quad (5.3)$$

235 
$$\int_0^{2\pi} \frac{(x-\xi)^2}{r^2} \sin \phi d\phi = \frac{\pi}{2} a \sin \theta, \quad (5.4)$$

236 
$$\int_0^{2\pi} \frac{(y-\eta)^2}{r^2} \cos \phi d\phi = \frac{\pi}{2} a \cos \theta, \quad (5.5)$$

237 
$$\int_0^{2\pi} \frac{(y-\eta)^2}{r^2} \sin \phi d\phi = -\frac{\pi}{2} a \sin \theta, \quad (5.6)$$

238 
$$\int_0^{2\pi} \frac{(x-\xi)(y-\eta)}{r^2} \cos \phi d\phi = -\frac{\pi}{2} a \sin \theta, \quad (5.7)$$

239 
$$\int_0^{2\pi} \frac{(x-\xi)(y-\eta)}{r^2} \sin \phi d\phi = -\frac{\pi}{2} a \cos \theta, \quad (5.8)$$

240 
$$\int_0^{2\pi} \frac{(x-\xi)^2}{r^2} \cos n\phi d\phi = \frac{\pi}{2} (1-a^2) \left(\frac{\rho}{R}\right)^{n-2} \cos(n-2)\theta, \quad n \geq 2, \quad (5.9)$$

241 
$$\int_0^{2\pi} \frac{(x-\xi)^2}{r^2} \sin n\phi d\phi = \frac{\pi}{2} (1-a^2) \left(\frac{\rho}{R}\right)^{n-2} \sin(n-2)\theta, \quad n \geq 2, \quad (5.10)$$

242 
$$\int_0^{2\pi} \frac{(y-\eta)^2}{r^2} \cos n\phi d\phi = -\frac{\pi}{2} (1-a^2) \left(\frac{\rho}{R}\right)^{n-2} \cos(n-2)\theta, \quad n \geq 2, \quad (5.11)$$



$$\int_0^{2\pi} \frac{(y - \eta)^2}{r^2} \sin n\phi d\phi = -\frac{\pi}{2}(1 - a^2) \left(\frac{\rho}{R}\right)^{n-2} \sin(n - 2)\theta, \quad n \geq 2, \quad (5.12)$$

$$\int_0^{2\pi} \frac{(x - \xi)(y - \eta)}{r^2} \cos n\phi d\phi = -\frac{\pi}{2}(1 - a^2) \left(\frac{\rho}{R}\right)^{n-2} \sin(n - 2)\theta, \quad n \geq 2, \quad (5.13)$$

$$\int_0^{2\pi} \frac{(x - \xi)(y - \eta)}{r^2} \sin n\phi d\phi = \frac{\pi}{2}(1 - a^2) \left(\frac{\rho}{R}\right)^{n-2} \cos(n - 2)\theta, \quad n \geq 2. \quad (5.14)$$

*Proof* We only prove a few of them, since the proof of the others is similar. First, since

$$\begin{aligned} (x - \xi)^2 &= (\rho \cos \theta - R \cos \phi)^2 = \rho^2 \cos^2 \theta + R^2 \cos^2 \phi - 2\rho R \cos \theta \cos \phi \\ &= \rho^2 \frac{1 + \cos 2\theta}{2} + R^2 \frac{1 + \cos 2\phi}{2} - 2\rho R \cos \theta \cos \phi, \end{aligned} \quad (5.15)$$

we obtain the integral on the left side of (5.1):

$$\begin{aligned} \int_0^{2\pi} \frac{(x - \xi)^2}{r^2} &= \frac{2\pi}{R^2(1 - a^2)} \left\{ \rho^2 \frac{1 + \cos 2\theta}{2} + R^2 \frac{1 + a^2 \cos 2\theta}{2} - 2\rho R a \cos^2 \theta \right\} \\ &= \frac{2\pi}{R^2(1 - a^2)} \left\{ \frac{R^2 + \rho^2}{2} - \rho^2 \right\} = \pi. \end{aligned} \quad (5.16)$$

Below we will show (5.9)–(5.16) first, and (5.3)–(5.8) afterwards. For (5.9), the integrand is given from (5.15)

$$\begin{aligned} (x - \xi)^2 \cos n\phi &= \rho^2 \left(\frac{1 + \cos 2\theta}{2}\right) \cos n\phi + R^2 \left(\frac{1 + \cos 2\phi}{2}\right) \cos n\phi \\ &\quad - 2\rho R \cos \theta \cos \phi \cos n\phi \\ &= \frac{\rho^2 + R^2}{2} \cos n\phi + \rho^2 \left(\frac{\cos 2\theta}{2}\right) \cos n\phi + \frac{R^2}{4} [\cos(n + 2)\phi \\ &\quad + \cos(n - 2)\phi] - \rho R \cos \theta [\cos(n + 1)\phi + \cos(n - 1)\phi]. \end{aligned} \quad (5.17)$$

262 Then the left side integral in (5.9) gives

$$\begin{aligned}
 & \int_0^{2\pi} \frac{(x - \xi)^2}{r^2} \cos n\phi d\phi \\
 &= \frac{2\pi}{R^2(1 - a^2)} \left\{ \frac{\rho^2 + R^2}{2} a^n \cos n\theta + \rho^2 \left( \frac{\cos 2\theta}{2} \right) a^n \cos n\theta \right. \\
 & \quad + \frac{R^2}{4} [a^{n+2} \cos(n + 2)\theta + a^{n-2} \cos(n - 2)\theta] \\
 & \quad \left. - \rho R \cos \theta [a^{n+1} \cos(n + 1)\theta + a^{n-1} \cos(n - 1)\theta] \right\} \\
 &= \frac{2\pi}{R^2(1 - a^2)} \left\{ \frac{\rho^2 + R^2}{2} a^n \cos n\theta + \frac{\rho^2}{4} a^n [\cos(n + 2)\theta + \cos(n - 2)\theta] \right. \\
 & \quad + \frac{R^2}{4} [a^{n+2} \cos(n + 2)\theta + a^{n-2} \cos(n - 2)\theta] \\
 & \quad \left. - \frac{\rho R}{2} a^{n+1} [\cos(n + 2)\theta + \cos n\theta] - \frac{\rho R}{2} a^{n-1} [\cos n\theta + \cos(n - 2)\theta] \right\}. \quad (5.18)
 \end{aligned}$$

270 In the above equations, the final coefficients in front of  $\cos n\theta$  and  $\cos(n + 2)\theta$  are  
 271 just zero. Hence we obtain

$$\int_0^{2\pi} \frac{(x - \xi)^2}{r^2} \cos n\phi d\phi = \frac{\pi a^{n-2}}{2R^2(1 - a^2)} \left[ R^2 + \frac{\rho^4}{R^2} - 2\rho^2 \right] \cos(n - 2)\theta. \quad (5.19)$$

273 There exists the equality,

$$\begin{aligned}
 R^2 + \frac{\rho^4}{R^2} - 2\rho^2 &= R^2 - \rho^2 + \rho^2 \left( \frac{\rho^2}{R^2} - 1 \right) \\
 &= (R^2 - \rho^2) \left\{ 1 - \frac{\rho^2}{R^2} \right\} = R^2(1 - a^2)^2. \quad (5.20)
 \end{aligned}$$

276 Combining (5.19) and (5.20) gives the desired result (5.9).

277 Next for (5.11), there exist the equalities,

$$\begin{aligned}
 (y - \eta)^2 &= (\rho \sin \theta - R \sin \phi)^2 = \rho^2 \sin^2 \theta + R^2 \sin^2 \phi - 2\rho R \sin \theta \sin \phi \\
 &= \rho^2 \frac{1 - \cos 2\theta}{2} + R^2 \frac{1 - \cos 2\phi}{2} - 2\rho R \sin \theta \sin \phi, \quad (5.21)
 \end{aligned}$$

280 and

$$\begin{aligned}
 (y - \eta)^2 \cos n\phi &= \frac{\rho^2 + R^2}{2} \cos n\phi - \frac{\rho^2}{2} \cos 2\theta \cos n\phi \\
 &\quad - \frac{R^2}{2} \cos 2\phi \cos n\phi - 2\rho R \sin \theta \sin \phi \cos n\phi \\
 &= \frac{\rho^2 + R^2}{2} \cos n\phi - \frac{\rho^2}{2} \cos 2\theta \cos n\phi \\
 &\quad - \frac{R^2}{4} [\cos(n + 2)\phi + \cos(n - 2)\phi] \\
 &\quad - \rho R \sin \theta [\sin(n + 1)\phi - \sin(n - 1)\phi]. \tag{5.22}
 \end{aligned}$$

286 Hence we have

$$\begin{aligned}
 \int_0^{2\pi} \frac{(y - \eta)^2}{r^2} \cos n\phi &= \frac{2\pi}{R^2(1 - a^2)} \left\{ \frac{\rho^2 + R^2}{2} a^n \cos n\theta - \frac{\rho^2}{2} \cos 2\theta a^n \cos n\theta \right. \\
 &\quad - \frac{R^2}{4} [a^{n+2} \cos(n + 2)\theta + a^{n-2} \cos(n - 2)\theta] \\
 &\quad \left. - \rho R \sin \theta [a^{n+1} \sin(n + 1)\theta - a^{n-1} \sin(n - 1)\theta] \right\} \\
 &= \frac{2\pi}{R^2(1 - a^2)} \left\{ \frac{\rho^2 + R^2}{2} a^n \cos n\theta - \frac{\rho^2}{4} a^n [\cos(n + 2)\theta \right. \\
 &\quad + \cos(n - 2)\theta] - \frac{R^2}{4} [a^{n+2} \cos(n + 2)\theta + a^{n-2} \cos(n - 2)\theta] \\
 &\quad - \frac{\rho R}{2} [a^{n+1} (\cos n\theta - \cos(n + 2)\theta) - a^{n-1} (\cos(n - 2)\theta \\
 &\quad \left. - \cos n\theta)] \right\} \\
 &= -\frac{\pi}{2(R^2 1 - a^2)} a^{n-2} \left( R^2 + \frac{\rho^4}{R^2} - 2\rho^2 \right) \cos(n - 2)\theta \\
 &= -\frac{\pi}{2} (1 - a^2) a^{n-2} \cos(n - 2)\theta, \tag{5.23}
 \end{aligned}$$

296 where we have used (5.20). This is the desired result (5.11).

297 Moreover for (5.14), we have

$$\begin{aligned}
 (x - \xi)(y - \eta) &= (\rho \cos \theta - R \cos \phi)(\rho \sin \theta - R \sin \phi) \\
 &= \rho^2 \cos \theta \sin \theta + R^2 \cos \phi \sin \phi - \rho R [\cos \theta \sin \phi + \sin \theta \cos \phi] \\
 &= \frac{\rho^2}{2} \sin 2\theta + \frac{R^2}{2} \sin 2\phi - \rho R [\cos \theta \sin \phi + \sin \theta \cos \phi],
 \end{aligned}$$

301 and

$$\begin{aligned}
 302 \quad (x - \xi)(y - \eta) \sin n\phi &= \frac{\rho^2}{2} \sin 2\theta \sin n\phi + \frac{R^2}{2} \sin 2\phi \sin n\phi \\
 303 \quad &\quad - \rho R [\cos \theta \sin \phi + \sin \theta \cos \phi] \sin n\phi \\
 304 \quad &= \frac{\rho^2}{2} \sin 2\theta \sin n\phi + \frac{R^2}{4} [-\cos(n+2)\phi + \cos(n-2)\phi] \\
 305 \quad &\quad - \frac{\rho R}{2} [\cos \theta (-\cos(n+1)\phi + \cos(n-1)\phi) \\
 306 \quad &\quad + \sin \theta (\sin(n+1)\phi + \sin(n-1)\phi)].
 \end{aligned}$$

307 Then we have

$$\begin{aligned}
 308 \quad \int_0^{2\pi} \frac{(x - \xi)(y - \eta)}{r^2} \sin n\phi &= \frac{2\pi}{R^2(1 - a^2)} \left\{ \frac{\rho^2}{2} \sin n\theta a^n \sin 2\theta \right. \\
 309 \quad &\quad + \frac{R^2}{4} [-a^{n+2} \cos(n+2)\theta + a^{n-2} \cos(n-2)\theta] \\
 310 \quad &\quad - \frac{\rho R}{2} [\cos \theta (-a^{n+1} \cos(n+1)\theta + a^{n-1} \cos(n-1)\theta) \\
 311 \quad &\quad \left. + \sin \theta (a^{n+1} \sin(n+1)\theta + a^{n-1} \sin(n-1)\theta)] \right\} \\
 312 \quad &= \frac{2\pi}{R^2(1 - a^2)} \left\{ \frac{\rho^2}{4} a^n [-\cos(n+2)\theta + \cos(n-2)\theta] \right. \\
 313 \quad &\quad + \frac{R^2}{4} [-a^{n+2} \cos(n+2)\theta + a^{n-2} \cos(n-2)\theta] \\
 314 \quad &\quad - \frac{\rho R}{4} \left\{ -a^{n+1} [\cos(n+2)\theta + \cos n\theta] + a^{n-1} [\cos n\theta + \cos(n-2)\theta] \right. \\
 315 \quad &\quad \left. + a^{n+1} [-\cos(n+2)\theta + \cos n\theta] + a^{n-1} [-\cos n\theta + \cos(n-2)\theta] \right\} \Big\} \\
 316 \quad &= \frac{\pi}{2R^2(1 - a^2)} \left( R^2 + \frac{\rho^4}{R^2} - 2\rho^2 \right) \left( \frac{\rho}{R} \right)^{n-2} \cos(n-2)\theta \\
 317 \quad &= \frac{\pi}{2} (1 - a^2) \left( \frac{\rho}{R} \right)^{n-2} \cos(n-2)\theta. \tag{5.24}
 \end{aligned}$$

318 This is the desired result (5.14).

319 Last, we show (5.3)–(5.8). Although we may follow the above approaches to derive  
 320 integrals directly, we will solicit different arguments by means of a remedy for (5.9)–  
 321 (5.14) with  $n = 1$ . First we have from (5.9) with  $n = 1$

$$322 \quad \int_0^{2\pi} \frac{(x - \xi)^2}{r^2} \cos \phi d\phi = \frac{\pi}{2} (1 - a^2) \left( \frac{\rho}{R} \right)^{-1} \cos(-\theta). \tag{5.25}$$

323 Let us scrutinize its derivation in (5.18). Note that in the expansions (3.18), there exist  
 324 no negative powers as  $(\frac{\rho}{R})^{-1}$ . In the last but one line in (5.18), we have found an  
 325 improper term

$$326 \quad \frac{2\pi}{R^2(1-a^2)} \frac{R^2 a^{-1}}{4} \cos \theta = \frac{\pi}{1-a^2} \frac{a^{-1}}{2} \cos \theta, \quad (5.26)$$

327 which should be replaced by the correct term

$$328 \quad \frac{\pi}{1-a^2} \frac{a}{2} \cos \theta. \quad (5.27)$$

329 Therefore, we find the error in (5.18) and so in (5.25) with  $n = 1$

$$\begin{aligned} 330 \quad E_1 &= \frac{\pi}{1-a^2} \frac{a}{2} \cos \theta - \frac{\pi}{1-a^2} \frac{a^{-1}}{2} \cos \theta \\ 331 \quad &= \frac{\pi}{2} \frac{1}{1-a^2} (a - a^{-1}) \cos \theta = \frac{\pi}{2} \frac{1}{1-a^2} a^{-1} (a^2 - 1) \cos \theta \\ 332 \quad &= -\frac{\pi}{2} a^{-1} \cos \theta. \end{aligned} \quad (5.28)$$

333 By removing this error, the true integral is obtained from (5.25) and (5.28)

$$\begin{aligned} 334 \quad \int_0^{2\pi} \frac{(x-\xi)^2}{r^2} \cos \phi d\phi &= \int_0^{2\pi} \frac{(x-\xi)^2}{r^2} \cos \phi d\phi + E_1 \\ 335 \quad &= \frac{\pi}{2} (1-a^2) \left(\frac{\rho}{R}\right)^{-1} \cos \theta - \frac{\pi}{2} a^{-1} \cos \theta \\ 336 \quad &= -\frac{\pi}{2} a \cos \theta. \end{aligned} \quad (5.29)$$

337 This is the desired result (5.3).

338 Finally, we show (5.8). From (5.14) with  $n = 1$ , we have

$$339 \quad \int_0^{2\pi} \frac{(x-\xi)(y-\eta)}{r^2} \sin n\phi d\phi = \frac{\pi}{2} (1-a^2) \left(\frac{\rho}{R}\right)^{-1} \cos \theta. \quad (5.30)$$

340 In its derivation (5.24) with  $n = 1$ , there also exists the error

$$341 \quad E_2 = \frac{\pi}{2} \frac{1}{1-a^2} (-a^{-1} + a) \cos \theta = -\frac{\pi}{2} a^{-1} \cos \theta. \quad (5.31)$$

342 By removing this error we have the true integral

$$\begin{aligned}
 343 \quad \int_0^{2\pi} \frac{(x-\xi)(y-\eta)}{r^2} \sin n\phi d\phi &= \int_0^{2\pi} \frac{(x-\xi)(y-\eta)}{r^2} \sin n\phi d\phi + E_2 \\
 344 \quad &= \frac{\pi}{2}(1-a^2) \left(\frac{\rho}{R}\right)^{-1} \cos\theta - \frac{\pi}{2} a^{-1} \cos\theta \\
 345 \quad &= -\frac{\pi}{2} a \cos\theta. \tag{5.32}
 \end{aligned}$$

346 This is the desired result (5.8). The proof of other formulas is similar, and this com-  
 347 pletes the proof of Lemma 5.1.

348 Note that for the integration computation of Fourier functions by means of orthog-  
 349 onality, we should also derive those formulas with  $n = 2$  due to  $\cos(n-2)\phi = 1$ .  
 350 However, since in (3.18), the constant is just  $\frac{1}{2}$ , fortunately, the final integral values in  
 351 (5.18) are exactly the same, so that we do not need extra-evaluation of (5.9). For the  
 352 same reason, we do not need an extra-evaluation for the expansions (4.17).

353 From Lemma 5.1 we have the Fourier expansions immediately.

354 **Theorem 5.1** Let  $\rho < R$ ,  $a = \frac{\rho}{R}$  and  $r^2 = R^2 + \rho^2 - 2\rho R \cos(\phi - \theta)$ . There exist  
 355 the expansions,

$$\begin{aligned}
 356 \quad \frac{(x-\xi)^2}{r^2} &= \frac{1}{2} - \frac{a}{2} \cos(\theta + \phi) + \frac{1-a^2}{2} \sum_{n=2}^{\infty} \left(\frac{\rho}{R}\right)^{n-2} \\
 357 \quad &\quad \times (\cos(n-2)\theta \cos n\phi + \sin(n-2)\theta \sin n\phi) \\
 358 \quad &= \frac{1}{2} - \frac{a}{2} \cos(\theta + \phi) + \frac{1-a^2}{2} \sum_{n=0}^{\infty} \left(\frac{\rho}{R}\right)^n \cos(n(\theta - \phi) - 2\phi), \\
 359 \quad \frac{(y-\eta)^2}{r^2} &= \frac{1}{2} + \frac{a}{2} \cos(\theta + \phi) \\
 360 \quad &\quad - \frac{1-a^2}{2} \sum_{n=2}^{\infty} \left(\frac{\rho}{R}\right)^{n-2} (\cos(n-2)\theta \cos n\phi + \sin(n-2)\theta \sin n\phi) \\
 361 \quad &= \frac{1}{2} + \frac{a}{2} \cos(\theta + \phi) - \frac{1-a^2}{2} \sum_{n=0}^{\infty} \left(\frac{\rho}{R}\right)^n \cos(n(\theta - \phi) - 2\phi), \\
 362 \quad \frac{(x-\xi)(y-\eta)}{r^2} &= -\frac{a}{2} \sin(\theta + \phi) + \frac{1-a^2}{2} \sum_{n=2}^{\infty} \left(\frac{\rho}{R}\right)^{n-2} \\
 363 \quad &\quad \times (-\sin(n-2)\theta \cos n\phi + \cos(n-2)\theta \sin n\phi) \\
 364 \quad &= -\frac{a}{2} \sin(\theta + \phi) - \frac{1-a^2}{2} \sum_{n=0}^{\infty} \left(\frac{\rho}{R}\right)^n \sin(n(\theta - \phi) - 2\phi).
 \end{aligned}$$

365 **6 The expansions of FS**

366 Based on Sects. 4 and 5, we provide new expansions of the FS for linear elastostatics  
 367 in 2D. For the simple FS with  $\rho < R$  we have their expansions from Theorem 4.1.

368 **Theorem 6.1** Let  $\rho < R$  and  $r^2 = R^2 + \rho^2 - 2\rho R \cos(\phi - \theta)$ . There exist the expan-  
 369 sions of the simple FS,

$$370 \quad \kappa \begin{pmatrix} \frac{x-\xi}{r^2} \\ \frac{y-\eta}{r^2} \end{pmatrix} = \frac{\kappa}{R} \sum_{n=0}^{\infty} \left(\frac{\rho}{R}\right)^n \begin{pmatrix} -\cos(n(\theta - \phi) - \phi) \\ \sin(n(\theta - \phi) - \phi) \end{pmatrix}.$$

371 Next for the principal FS with  $\rho < R$ , we have their expansions from Theorem 5.1.

372 **Theorem 6.2** Let  $\rho < R$ ,  $a = \frac{\rho}{R}$  and  $r^2 = R^2 + \rho^2 - 2\rho R \cos(\phi - \theta)$ . There exist  
 373 the expansions of the principal FS,

$$374 \quad E_2^i(\mathbf{x} - \xi) = \begin{pmatrix} T_{11}^i & T_{12}^i \\ T_{12}^i & T_{22}^i \end{pmatrix},$$

375 where the entries

$$376 \quad T_{11}^i = (1 - \kappa) \left\{ - \left[ \ln R - \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{\rho}{R}\right)^n \cos n(\theta - \phi) \right] \right. \\
 377 \quad \left. + D \left[ \frac{1}{2} - \frac{a}{2} \cos(\theta + \phi) + \frac{1 - a^2}{2} \sum_{n=0}^{\infty} \left(\frac{\rho}{R}\right)^n \cos(n(\theta - \phi) - 2\phi) \right] \right\},$$

$$378 \quad T_{22}^i = (1 - \kappa) \left\{ - \left[ \ln R - \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{\rho}{R}\right)^n \cos n(\theta - \phi) \right] \right. \\
 379 \quad \left. + D \left[ \frac{1}{2} + \frac{a}{2} \cos(\theta + \phi) - \frac{1 - a^2}{2} \sum_{n=0}^{\infty} \left(\frac{\rho}{R}\right)^n \cos(n(\theta - \phi) - 2\phi) \right] \right\},$$

$$380 \quad T_{12}^i = (1 - \kappa) D \left[ -\frac{a}{2} \sin(\theta + \phi) - \frac{1 - a^2}{2} \sum_{n=0}^{\infty} \left(\frac{\rho}{R}\right)^n \sin(n(\theta - \phi) - 2\phi) \right].$$

381 In the NFM, we do need the Fourier expansions for both  $\rho > R$  and  $\rho < R$ . We  
 382 have the following theorem.

383 **Theorem 6.3** Let  $\rho > R$  and  $r^2 = R^2 + \rho^2 - 2\rho R \cos(\phi - \theta)$ . There exist the  
 384 expansions of the simple FS,

$$385 \quad \kappa \begin{pmatrix} \frac{x-\xi}{r^2} \\ \frac{y-\eta}{r^2} \end{pmatrix} = \frac{\kappa}{\rho} \sum_{n=0}^{\infty} \left(\frac{R}{\rho}\right)^n \begin{pmatrix} \cos(n(\phi - \theta) - \theta) \\ -\sin(n(\phi - \theta) - \theta) \end{pmatrix}. \quad (6.1)$$

386 *Proof* We will use Theorem 6.1 via symmetry. When  $\rho > R$ , we have

$$387 \quad \frac{x - \xi}{r^2} = \frac{\rho \cos \theta - R \cos \phi}{R^2 + \rho^2 - 2\rho R \cos(\phi - \theta)} = -\frac{R \cos \phi - \rho \cos \theta}{R^2 + \rho^2 - 2\rho R \cos(\theta - \phi)}. \quad (6.2)$$

388 If switching  $(\rho, \theta)$  and  $(R, \phi)$ , from Theorem 6.1 and (6.2), we obtain the first com-  
389 ponent of the vector in (6.1):

$$390 \quad \kappa \frac{x - \xi}{r^2} = \frac{\kappa}{\rho} \sum_{n=0}^{\infty} \left(\frac{R}{\rho}\right)^n \cos(n(\phi - \theta) - \theta). \quad (6.3)$$

391 The proof for the second component of the vector in (6.1) is similar, and this completes  
392 the proof of Theorem 6.3.

393 Similarly, we have the following expansions from Theorem 6.2 via the symmetry.

394 **Theorem 6.4** Let  $\rho > R$ ,  $a = \frac{\rho}{R}$  and  $r^2 = R^2 + \rho^2 - 2\rho R \cos(\phi - \theta)$ . There exist  
395 the expansions of the principal FS,

$$396 \quad E_2^e(\mathbf{x} - \xi) = \begin{pmatrix} T_{11}^e & T_{12}^e \\ T_{12}^e & T_{22}^e \end{pmatrix},$$

397 where the entries are given by

$$398 \quad T_{11}^e = (1 - \kappa) \left\{ - \left[ \ln \rho - \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{R}{\rho}\right)^n \cos n(\theta - \phi) \right] \right. \\ 399 \quad \left. + D \left[ \frac{1}{2} - \frac{a^{-1}}{2} \cos(\theta + \phi) + \frac{1 - a^{-2}}{2} \sum_{n=0}^{\infty} \left(\frac{R}{\rho}\right)^n \cos(n(\phi - \theta) - 2\theta) \right] \right\},$$

$$400 \quad T_{22}^e = (1 - \kappa) \left\{ - \left[ \ln \rho - \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{R}{\rho}\right)^n \cos n(\theta - \phi) \right] \right. \\ 401 \quad \left. + D \left[ \frac{1}{2} + \frac{a^{-1}}{2} \cos(\theta + \phi) - \frac{1 - a^{-2}}{2} \sum_{n=0}^{\infty} \left(\frac{R}{\rho}\right)^n \cos(n(\phi - \theta) - 2\theta) \right] \right\},$$

$$402 \quad T_{12}^e = (1 - \kappa) D \left[ -\frac{a^{-1}}{2} \sin(\theta + \phi) - \frac{1 - a^{-2}}{2} \sum_{n=0}^{\infty} \left(\frac{R}{\rho}\right)^n \sin(n(\phi - \theta) - 2\theta) \right].$$

403 All series expansions in this paper have been verified by computation.

404 *Remark 6.1* The principal FS is more important than the simple FS, because from our  
405 numerical experiments, we may not need the simple FS (see [13]). By series opera-  
406 tions, some expansions for  $\frac{(x-\xi)^2}{r^2}$ ,  $\frac{(y-\eta)^2}{r^2}$  and  $\frac{(x-\xi)(y-\eta)}{r^2}$  are given in Chen et al. [5],  
407 which are more complicated than those in Theorem 5.1. The simplicity of expansions  
408 of FS in this paper is imperative to real application of linear elastostatics, such as error  
409 and stability analysis of MFS in [14, 15], and algorithms and analysis of the NFM  
410 in [17].



411 **References**

- 412 1. Chen G, Zhou J (1992) Boundary element methods. Academic Press, New York
- 413 2. Chen JT, Kuo SR, Lin JH (2002) Analytical study and numerical experiments for degenerate scale
- 414 problems in the boundary element method for two-dimensional elasticity. *Eng Anal Boundary*
- 415 *Elements* 26:1669–1681
- 416 3. Chen JT, Shen WC, Wu AC (2006) Null-field integral equations for stress field around circular holes
- 417 under antiplane shear. *Eng Anal Boundary Elements* 30:205–217
- 418 4. Chen JT, Lee CF, Chen JL, Lin JH (2002) An alternative method for degenerate scale problem in bound-
- 419 ary element methods for two-dimensional Laplace equation. *Eng Anal Boundary Elements* 26:559–569
- 420 5. Chen JT, Lee YT, Chou KH (2010) Revisit of two classical elasticity problems by using the null-fields
- 421 integration equations. *J Mech* 26(3):113–121
- 422 6. Chen JT, Lee YT, Lee JW (2010) Torsional rigidity of an elliptic bar with multiple elliptic inclusions
- 423 using a null-field integral approach. *Comput Mech* 46:511–519
- 424 7. Chen JT, Lee JW, Lee SY (2010) Analytical and numerical investigation for true and spurious eigen-
- 425 solutions of an elliptical membrane using the real-part dual BEM, Technical report
- 426 8. Crouch SL, Mogilevskaya SG (2003) On the use of Somigliana's formula and Fourier series for elas-
- 427 ticity problems with circular boundaries. *Int J Numer Meth Eng* 58:537–578
- 428 9. Gradsheyan IS, Ryzhik ZM (1965) Tables of integrals, series and products. Academic Press, New York
- 429 10. Hsiao GC, Wendland WL (2008) Boundary integral equations. Springer, Berlin
- 430 11. Li ZC (1998) Combined methods for elliptic equations with singularities, interfaces and infinities.
- 431 Kluwer Academic Publishers, Boston
- 432 12. Li ZC (2009) Method of fundamental solutions for annular shaped domains. *J Comput Appl Math*
- 433 228:355–372
- 434 13. Li ZC (2010) Error analysis and numerical computation for hybrid Trefftz methods coupling traction
- 435 conditions in linear elastostatics, A comparison, revised version for Numer. PDE
- 436 14. Li ZC (2010) Error analysis of the method of fundamental solutions for linear elastostatics, Technical
- 437 report
- 438 15. Li ZC, Huang J, Huang HT (2010) Stability analysis of method of fundamental solutions for mixed
- 439 boundary value problems of Laplace's equation. *Computing* 88:1–29
- 440 16. Li ZC et al (2010) The explicit algorithms of the null-field method for Laplace's equation on circular
- 441 domains with multiple holes, Technical report, National Sun Yat-sen University, Kaohsiung, Taiwan
- 442 17. Li ZC et al (2010) The explicit algorithms of the null-field method for linear elastostatics on circular
- 443 domains with multiple holes, Technical report, National Sun Yat-sen University, Kaohsiung, Taiwan
- 444 18. Li ZC, Lee MG, Chen JT (2010) The error analysis for the null field method for Laplace's equation in
- 445 circular domains with multiple holes, Technical report, National Sun Yat-sen University, Kaohsiung,
- 446 Taiwan
- 447 19. Li ZC, Lu TT, Hu HY, Cheng AHD (2008) Trefftz and collocation methods. WIT press, Southampton
- 448 20. Mogilevskaya SG, Linkov AM (1998) Complex fundamental solutions and complex variables bound-
- 449 ary element method in elasticity. *Comput Mech* 22:88–92
- 450 21. Qin QH (2000) The Trefftz finite and boundary element method. WIT press, Southampton, pp 282
- 451 22. Yu DH (2002) Natural boundary integral method and its applications. Science Press/Kluwer Academic
- 452 Publishers, Beijing/New York